

WASTEWATER-BASED EPIDEMIOLOGICAL SURVEILLANCE OF THE SARS-COV-2

— WEEKLY REPORT

— RESULTS OF 13/09/2023

Authors: R. Janssens¹, H. Maloux¹, S. Hanoteaux¹, V. Hutse¹, L. Van Poelvoorde², N. Roosens², B. Verhaegen³, K. Van Hoorde³, K. Dierick³, K. Blot¹, M. Lesenfants^{1*}

¹ Sciensano, Service Epidemiology of infectious diseases

² Sciensano, Transversal activities in Applied Genomics

³ Sciensano, Service Foodborne pathogens

*Contact and Info: wastewater.info@sciensano.be

In collaboration with:



1. Table of contents

1. Table of contents	2
2. Summary	3
3. Introduction	4
4. Methodology	4
4.1. SAMPLE COLLECTION AND ANALYSIS	4
4.2. WASTEWATER RESULTS	5
4.3. ALERTING INDICATORS	5
4.4. CAUTION POINTS FOR THE RESULTS INTERPRETATION.....	6
5. Results	7
5.1. NATIONAL LEVEL	7
5.2. REGIONAL LEVEL.....	10
5.3. PROVINCIAL LEVEL	12
5.4. INDIVIDUAL CATCHMENT AREAS LEVEL.....	13
6. Appendix – Areas classified by alerting indicator	15

2. Summary

In the present work, the circulation of the SARS-CoV-2 virus in the environment is assessed based on three alerting indicators. The analysis of wastewater samples collected in 41 wastewater treatment plants covers 45% of the Belgian population. The results of the wastewater surveillance are a complementary source of information to the infection cases number. Indeed, the wastewater results do notably include all asymptomatic persons, and are independent of the testing strategy.

Here are the conclusions based on the latest results of September 13th 2023 (Week 37)¹:

- At the national level: A maximum number of areas in High Circulation was recorded on the 13th September 2023 after several weeks of increase. Since this date, the High Circulation indicator is decreasing with 7 areas positive this week. Overall, the viral loads are at a high level and are decreasing.
 - At the regional level: The viral loads are high and increasing in Wallonia.
 - At the provincial level: The Increasing Trend indicator is positive in 3 provinces, and was in 6 last week. The High Circulation indicator is positive in 3 provinces, and was in 5 last week.
 - At the covered areas level: among the 34 areas covered, the number of treatment plants positive to the different indicators are 12 for the Increasing Trend indicator, 3 for the Fast Increase indicator, and 7 for the High Circulation indicator.
-

The wastewater situation can be followed on a weekly basis on:

- The graphics available on the public [COVID-19 dashboard](#)
 - The epidemiological update published on Fridays by the Risk Assessment Group (RAG) after validation by the Risk Management Group (RMG) in [French](#) and [Dutch](#).
 - Further details on the methodology applied for the wastewater surveillance can be found in [the Appendix Methodology document \(access available online\)](#).
-

¹ The wastewater surveillance is based on 41 areas. However, due to technical issues, the area of Amay, Brussels-North, Destelbergen, Houthalen-Centrum, Liege (Grosses Battes), Liege Oupeye, and Liege Sclessin were not sampled this week (see Appendix A5). Hence, this week's report is only based on 34 areas.

3. Introduction

The national wastewater-based surveillance of COVID-19 started in September 2020. The present report aims to assess the wastewater-based epidemiological situation in Belgium. It is updated weekly on Tuesday based on the concentration measured in the samples during the previous week.

The SARS-CoV-2 concentrations are measured twice per week in 41 wastewater treatment plants (WWTPs). The evolution of these concentrations is assessed in the present report thanks to three wastewater-based alerting indicators. The assessment is performed at 4 different spatial levels: national, regional, provincial, and the areas covered by the treatment plants.

Finally, the remaining sources of uncertainties are discussed together with their expected impacts on the interpretation of the wastewater results.

4. Methodology

4.1. SAMPLE COLLECTION AND ANALYSIS

Samples are collected twice a week in several wastewater treatment plants (WWTP) covering around 42% of the Belgian population: 36% in the Flemish region, 34.5% in the Walloon region, and nearly 100% in the Brussels region. Figure 1 shows the catchment areas covered by the WWTP located in areas with high population density. The catchment area of a WWTP corresponds to the geographical area from which the wastewater are collected.

Samples are collected on Mondays and Wednesdays by auto-samplers (24-hour composite) at the influent of WWTP. The analysis are distributed amongst 3 different laboratories to quantify the concentration of SARS-CoV-2 RNA. The results corresponding to the samples of the Mondays and Wednesdays are made publicly available on Wednesdays and Fridays, respectively.

Further details on the coverage, sampling plan, and analytical method can be found in [the Appendix Methodology document](#)




Figure 1: The population located in the areas covered by the wastewater treatment plants (highlighted in yellow) and the population density for each municipality (indicated by the blue scale).

4.2. WASTEWATER RESULTS

In order to account for possible dilution by rainy events and for the number of people living in the catchment area of each WWTP, a correction is applied on the viral concentration: the concentration is multiplied by the inlet flow and divided by the number of inhabitant of its respective WWTP and expressed by 100k inhabitant. The viral loads are expressed in copy/day/100k inhabitants.

The limit of quantification of the analytical method was estimated at 10 copies/ml.

4.3. ALERTING INDICATORS

To highlight the areas of possible concern, the three alerting indicators are assessed twice a week, based on viral loads (RNA copies/day/100k inhabitants):

1. The **Increasing Trend** indicator highlights the catchment areas where the viral loads have been increasing for more than 6 days. The indicator is computed based on the moving average on the past two weeks of the viral load.
2. The **Fast Increase** indicator highlights the catchment areas where the viral loads have rapidly increased for the last week. It corresponds to a situation where the moving average on the past 7 days of the viral load has increased faster than 70% per week if being above the estimated limit of quantification. The increasing slope is normalized for each treatment plant.
3. The **High Circulation** indicator highlights the catchment areas where the viral loads are high. It corresponds to a situation where the viral loads exceed half of the highest value recorded during the ninth wave (i.e. from 21th of November 2022 till first of January 2023).

The indicators were developed to monitor the different phases of an outbreak. the indicator Increasing Trend will be on alert first in the emerging phase. If the viral loads increase quickly, the Fast Increase indicator will then turn on alert. Finally, the viral loads may be sufficiently high to result in the High Circulation indicator to turn on alert.

4.4. CAUTION POINTS FOR THE RESULTS INTERPRETATION

The viral loads should not be directly compared between catchment areas, provinces, nor regions because analyses are conducted by three different laboratories. To mitigate this bias, the indicators are computed on normalized viral loads, allowing for comparison between the different areas. Further details are available in [the methodology document \(access online\)](#).

5. Results

5.1. NATIONAL LEVEL

Table 1 shows, at the national level, the results obtained on the last sample of September 13th 2023, week 37, compared to the ones obtained on September 06th 2023. In this table, any change in indicator positivity (i.e. if the value for any indicator has changed from 0 to 1 or from 1 to 0) is indicated in **coloured bold text**.

None of the three indicators is positive this week.

Table 1: Alerting indicators positive (1) or not (0) on September 13th 2023. Columns represent the population coverage of Belgium (Pop. coverage) and the three alerting indicators High Circulation (High), Fast Increase (Fast) and Increasing Trend (Incr.). The specifications of the four last columns are explained in the footnotes 1-4 below the table. Missing data is indicated with a “/”.

Country	Pop. coverage	High	Fast	Incr.	Norm. viral load (%) ¹	Mean viral load ²	Norm. evol. (%/week) ³	Incr. days ⁴
Belgium	44%	0	0	0	23	0.29	-7	4

¹ : the viral load normalized with the maximum viral load measured in the corresponding catchment area during the ninth wave (i.e. from 21th of November 2022 till the first of January 2023).

² : the viral load computed on the mean of the replicate of the three targeted gene fragments as explained in section “3.2 Wastewater results”. The mean viral load is expressed in 10¹² copies/day/100k inhabitants.

³ : the slope (%/week) of the past 7 days moving average of the viral load if the corresponding concentration is above the estimated limit of quantification.

⁴ : the cumulative number of days of increase of the past 14 days moving average of the viral load.

The geographical location of the areas is presented in Figure 2 together with the status of the three indicators.

A maximum number of areas in High Circulation was recorded on the 13th September 2023 after several weeks of increase. Since this date, the High Circulation indicator is decreasing with 7 areas positive this week. Overall, the viral loads are at a high level and are decreasing.




Figure 2: The geographical location of the covered areas with the corresponding status of the three alerting indicators: High circulation, Fast Increase and Increasing trend. If an indicator is positive its corresponding slice is displayed in its colour (see legend) whereas when negative the same slice is greyed out. The names of the covered areas with respect of their localization can be found in Figure 1.



Figure 3: Number of areas having a positive alerting indicator (coloured bars), number of areas having a negative indicator (lighter coloured bars) and number of areas not sampled (greyed out bars). Latest results correspond to September 13th 2023, week 37.

5.2. REGIONAL LEVEL

Figure 4 shows, at the regional level, the viral loads in the wastewaters.

Two waves can be seen in Figure 4:

- The 9th wave starting on 21st November 2022.
- The 10th wave starting on 23rd January 2023.




Figure 4: The SARS-CoV-2 RNA viral loads expressed as copies/days/100k inhabitants (based on the past two weeks moving average) for the three regions.

Table 2 shows, at the regional level, the results obtained on the last sample of Wednesday September 13th 2023, week 37. It allows to track the changes between the situation as of today (September 13th 2023) and the situation as of last week (September 06th 2023). Hereby, two distinct cases are taken into account:

1. If a region has at least one indicator fulfilled this week and it was not the case last week, its **name is displayed in bold** in the table.
2. If a region has at least one indicator fulfilled this week and if it also was the case last week, any change in indicator fulfilment (i.e. if the value for any indicator has changed from 0 to 1 or from 1 to 0) is indicated in **coloured bold text**.

The High Circulation and Increasing Trend indicators are positive in Wallonia.

Table 2: Alerting indicators positive (1) or not (0) on September 13th 2023. Columns represent the population coverage of the regions (Pop. coverage) and the three alerting indicators High Circulation (High), Fast Increase (Fast) and Increasing Trend (Incr.). The specifications of the four last columns are explained in the footnotes 1-4 below the table. Missing data is indicated with a “/”.

Region	Pop. coverage	High	Fast	Incr.	Norm. viral load (%) ¹	Mean viral load ²	Norm. evol. (%/week) ³	Incr. days ⁴
Brussels	100%	0	0	0	6.4	0.45	-29	0
Flanders	41%	0	0	0	12.8	0.19	0	4
Wallonia	31%	1	0	1	54.6	0.54	4	7

¹ : the viral load normalized with the maximum viral load measured in the corresponding catchment area during the ninth wave (i.e. from 21th of November 2022 till the first of January 2023).

² : the viral load computed on the mean of the replicate of the three targeted gene fragments as explained in section “3.2 Wastewater results”. The mean viral load is expressed in 10¹² copies/day/100k inhabitants.

³ : the slope (%/week) of the past 7 days moving average of the viral load if the corresponding concentration is above the estimated limit of quantification.

⁴ : the cumulative number of days of increase of the past 14 days moving average of the viral load.

5.3. PROVINCIAL LEVEL

Table 3 shows, at the provincial level, the results obtained on the last sample of Wednesday September 13th 2023, week 37. It allows to track the changes between the situation as of today (September 13th 2023) and the situation as of last week (September 06th 2023). Hereby, two distinct cases are taken into account:

1. If a province has at least one indicator fulfilled this week and it was not the case last week, its **name is displayed in bold** in the table.
2. If a province has at least one indicator fulfilled this week and if it also was the case last week, any change in indicator fulfilment (i.e. if the value for any indicator has changed from 0 to 1 or from 1 to 0) is indicated in **coloured bold text**.

Table 3 shows, for each Province, the results associated with the samples of Wednesday September 13th 2023, for the three alerting indicators:

- The Increasing Trend indicator is positive in 3 provinces: Limburg, Namur and Oost-Vlaanderen, and was in 6 last week.
- The Fast Increase indicator is only positive in the province of Luxembourg, and was in 3 last week.
- The High Circulation indicator is positive in 3 provinces: Brabant Wallon, Luxembourg and Namur, and was in 5 last week.

Table 3: Alerting indicators positive (1) or not (0) on September 13th 2023. Columns represent the population coverage of the WWTPs within the Province (Pop. coverage) and the three alerting indicators High Circulation (High), Fast Increase (Fast) and Increasing Trend (Incr.). The specifications of the four last columns are explained in the footnotes 1-4 below the table. Missing data is indicated with a “/”.

Province	Pop. coverage	High	Fast	Incr.	Norm. viral load (%) ¹	Mean viral load ²	Norm. evol. (%/w) ³	Incr. days ⁴
Antwerpen	3580%	0	0	0	30.8	0.22	0	0
Brabant Wallon	4400%	1	0	0	70.3	0.53	-26	5
Brussels	10000%	0	0	0	6.4	0.45	-29	0
Hainaut	3380%	0	0	0	29.0	0.35	2	5
Liège	4240%	/	/	/	/	/	/	/
Limburg	2380%	0	0	1	8.3	0.10	0	9
Luxembourg	980%	1	1	0	125.2	1.76	155	2
Namur	2530%	1	0	1	115.0	1.01	25	20
Oost-Vlaanderen	3420%	0	0	1	6.9	0.20	0	7
Vlaams-Brabant	4320%	0	0	0	2.6	0.18	0	6
West-Vlaanderen	4090%	0	0	0	3.7	0.17	0	2

¹ : the viral load normalized with the maximum viral load measured in the corresponding catchment area during the ninth wave (i.e. from 21th of November 2022 till the first of January 2023).

² : the viral load computed on the mean of the replicates of the three targeted gene fragments as explained in section “3.2 Wastewater results”. The mean viral load is expressed in 10¹² copies/day/100k inhabitants.

³ : the slope (%/week) of the past 7 days moving average of the viral load if the corresponding concentration is above the estimated limit of quantification.

⁴ : the cumulative number of days of increase of the past 14 days moving average of the viral load.

5.4. INDIVIDUAL CATCHMENT AREAS LEVEL

Table 4 shows, at the catchment area level, the results obtained on the last sample of Wednesday September 13th 2023, week 37. It allows to track the changes between the situation as of today (September 13th 2023) and the situation as of last week (September 06th 2023). Hereby, three distinct cases are taken into account:

1. If an area has at least one indicator fulfilled this week and it was not the case last week, its **name is displayed in bold** in the table.
2. If an area has at least one indicator fulfilled this week and if it also was the case last week, any change in indicator fulfilment (i.e. if the value for any indicator has changed from 0 to 1 or from 1 to 0) is indicated in **coloured bold text**.

Any area which had at least one indicator fulfilled last week but not this week is listed below Table 4.

Here are the results associated with the samples of September 13th 2023:

- The Increasing Trend indicator is positive in 12 covered areas. Amongst these areas, the viral load is continually increasing since two or more weeks in at least one area. Further details can be found in Appendix A3. Last week, this indicator was positive in 24 covered areas.
- The Fast Increase is positive in 3 covered areas: Arlon (166.6% increase per week), Marche-en-Famenne (128.6% increase per week) and Mornimont (87.2% increase per week) (see Appendix A2 for more details). Last week, this indicator was positive in 13 covered areas.
- The High Circulation indicator is positive in 7 covered areas: Marche-en-Famenne (159.34%), Namur-Brumagne (121.7%), Rosieres (Lasne) (114.31%), Arlon (110.03%), Mornimont (98.89%), Vallee du Hain (L'Orchis) (75.42%) and Antwerpen-Zuid (57.77%) (see Appendix A1 for more details). Last week, this indicator was positive in 14 covered areas.
- An alerting situation is evidenced for the covered areas of Marche-en-Famenne and Mornimont as all three indicators are positive.

The wastewater results at the level of the local covered areas can be accessed online for each area on the [COVID-19 dashboard](#).

Table 4: Alerting indicators positive (1) or not (0) on September 13th 2023. Columns represent the provinces, different WWTPs within the Provinces and the three alerting indicators High Circulation (High), Fast Increase (Fast) and Increasing Trend (Incr.). The specifications of the four last columns are explained in the footnotes 1-4 below the table. Missing data is indicated with a “/”.

Province	WWTP	High	Fast	Incr.	Norm. viral load (%) ¹	Mean viral load ²	Norm evol. (%/week) ³	Incr. days ⁴
Antwerpen	Antwerpen-Zuid	1	0	0	57.77	0.47	-19.1	0
Luxembourg	Arlon	1	1	0	110.03	1.52	166.6	0
Oost-Vlaanderen	Dendermonde	0	0	1	10.00	0.05	0.0	14
Limburg	Genk	0	0	1	12.83	0.19	0.0	14
Oost-Vlaanderen	Gent	0	0	1	5.71	0.26	0.0	9
Vlaams-Brabant	Leuven	0	0	1	2.22	0.17	0.0	7
Vlaams-Brabant	Liedekerke	0	0	1	2.74	0.14	0.0	14

Province	WWTP	High	Fast	Incr.	Norm. viral load (%) ¹	Mean viral load ²	Norm evol. (%/week) ³	Incr. days ⁴
Luxembourg	Marche-en-Famenne	1	1	1	159.34	2.31	128.6	7
Namur	Mornimont	1	1	1	98.89	2.06	87.2	35
Namur	Namur-Brumagne	1	0	1	121.70	0.57	-0.3	14
West-Vlaanderen	Roeselare	0	0	1	8.44	0.28	-25.3	16
Brabant Wallon	Rosieres (Lasne)	1	0	1	114.31	0.90	-34.8	16
Limburg	Tessengerlo	0	0	1	14.14	0.11	0.0	16
Brabant Wallon	Vallee du Hain (L'Orchis)	1	0	0	75.42	0.59	0.0	2
Hainaut	Wasmuel	0	0	1	40.87	0.48	43.6	14

¹ : the viral load normalized with the maximum viral load measured in the corresponding catchment area during the ninth wave (i.e. from 21th of November 2022 till the first of January 2023).

² : the viral load computed on the mean of the replicates of the three targeted gene fragments as explained in section "3.2 Wastewater results"; The mean viral load is expressed in 10¹² copies/day/100k inhabitants.

³ : the slope (%/week) of the past 7 days moving average of the viral load if the corresponding concentration is above the estimated limit of quantification.

⁴ : the cumulative number of days of increase of the past 14 days moving average of the viral load.

The following areas had the corresponding indicator positive last week but not this week:

- Increasing Trend indicator: Aalst, Aartselaar, Antwerpen-Noord, Beersel, Brugge, Brussels-South, Deurne, Grimbergen, Harelbeke, Hasselt, Marchienne-au-Pont, Mechelen-Noord and Montignies-sur-Sambre.
- Fast Increase indicator: Brugge, Deurne, Marchienne-au-Pont and Montignies-sur-Sambre.
- High Circulation indicator: Antwerpen-Noord, Basse Wavre (Dyle), Deurne, Froyennes, Marchienne-au-Pont, Montignies-sur-Sambre and Roselies.

Further details on covered area without positive indicators can be found in Table A4.

6. Appendix – Areas classified by alerting indicator

Table A1: Areas for which the High Circulation indicator is positive (7 out of 41 on September 13th 2023).

Province	WWTP	High	Fast	Incr.	Norm. viral load (%) ¹	Mean viral load ²	Norm evol. (%/week) ³	Incr. days ⁴	Date Max cc ⁵
Luxembourg	Marche-en-Famenne	1	1	1	159.34	2.31	128.6	7	13/09/2023
Namur	Namur-Brumagne	1	0	1	121.70	0.57	-0.3	14	11/09/2023
Brabant Wallon	Rosieres (Lasne)	1	0	1	114.31	0.90	-34.8	16	06/09/2023
Luxembourg	Arlon	1	1	0	110.03	1.52	166.6	0	30/08/2023
Namur	Mornimont	1	1	1	98.89	2.06	87.2	35	9 th wave
Brabant Wallon	Vallee du Hain (L'Orchis)	1	0	0	75.42	0.59	0.0	2	21/08/2023
Antwerpen	Antwerpen-Zuid	1	0	0	57.77	0.47	-19.1	0	15/03/2023

¹ : the viral load normalized with the maximum viral load measured in the corresponding catchment area during the ninth wave (i.e. from 21th of November 2022 till the first of January 2023).

² : the viral load computed on the mean of the replicates of the three targeted gene fragments as explained in section "3.2 Wastewater results". The mean viral load is expressed in 10¹² copies/day/100k inhabitants.

³ : the slope (%/week) of the past 7 days moving average of the viral load if the corresponding concentration is above the estimated limit of quantification.

⁴ : the cumulative number of days of increase of the past 14 days moving average of the viral load.

⁵ : date at which the measured viral load was the highest since the beginning of the ninth wave. If the date was between the 21th of November 2022 and the first of January 2023, the date is considered to be during the ninth wave and mentioned as such.

Table A2: Areas for which the Fast Increase indicator is positive (3 out of 41 on the September 13th 2023).

Province	WWTP	High	Fast	Incr.	Norm. viral load (%) ¹	Mean viral load ²	Norm evol. (%/week) ³	Incr. days ⁴	Date Max cc ⁵
Luxembourg	Arlon	1	1	0	110.03	1.52	166.6	0	30/08/2023
Luxembourg	Marche-en-Famenne	1	1	1	159.34	2.31	128.6	7	13/09/2023
Namur	Mornimont	1	1	1	98.89	2.06	87.2	35	9 th wave

¹ : the viral load normalized with the maximum viral load measured in the corresponding catchment area during the ninth wave (i.e. from 21th of November 2022 till the first of January 2023).

² : the viral load computed on the mean of the replicates of the three targeted gene fragments as explained in section "3.2 Wastewater results". The mean viral load is expressed in 10¹² copies/day/100k inhabitants.

³ : the slope (%/week) of the past 7 days moving average of the viral load if the corresponding concentration is above the estimated limit of quantification.

⁴ : the cumulative number of days of increase of the past 14 days moving average of the viral load.

⁵ : date at which the measured viral load was the highest since the beginning of the ninth wave. If the date was between the 21th of November 2022 and the first of January 2023, the date is considered to be during the ninth wave and mentioned as such.

Table A3: Areas for which the Increasing Trend indicator is positive (12 out of 41 on the September 13th 2023).

Province	WWTP	High	Fast	Incr.	Norm. viral load (%) ¹	Mean viral load ²	Norm evol. (%/week) ³	Incr. days ⁴	Date Max cc ⁵
Namur	Mornimont	1	1	1	98.89	2.06	87.2	35	9 th wave
West-Vlaanderen	Roeselare	0	0	1	8.44	0.28	-25.3	16	05/10/2022
Brabant Wallon	Rosieres (Lasne)	1	0	1	114.31	0.90	-34.8	16	06/09/2023
Limburg	Tessenderlo	0	0	1	14.14	0.11	0.0	16	15/03/2023
Oost-Vlaanderen	Dendermonde	0	0	1	10.00	0.05	0.0	14	06/02/2023
Limburg	Genk	0	0	1	12.83	0.19	0.0	14	08/03/2023
Vlaams-Brabant	Liedekerke	0	0	1	2.74	0.14	0.0	14	9 th wave
Namur	Namur-Brumagne	1	0	1	121.70	0.57	-0.3	14	11/09/2023
Hainaut	Wasmuel	0	0	1	40.87	0.48	43.6	14	9 th wave
Oost-Vlaanderen	Gent	0	0	1	5.71	0.26	0.0	9	08/03/2023
Vlaams-Brabant	Leuven	0	0	1	2.22	0.17	0.0	7	9 th wave
Luxembourg	Marche-en-Famenne	1	1	1	159.34	2.31	128.6	7	13/09/2023

¹ : the viral load normalized with the maximum viral load measured in the corresponding catchment area during the ninth wave (i.e. from 21th of November 2022 till the first of January 2023).

² : the viral load computed on the mean of the replicates of the three targeted gene fragments as explained in section "3.2 Wastewater results". The mean viral load is expressed in 10¹² copies/day/100k inhabitants.

³ : the slope (%/week) of the past 7 days moving average of the viral load if the corresponding concentration is above the estimated limit of quantification.

⁴ : the cumulative number of days of increase of the past 14 days moving average of the viral load.

⁵ : date at which the measured viral load was the highest since the beginning of the ninth wave. If the date was between the 21th of November 2022 and the first of January 2023, the date is considered to be during the ninth wave and mentioned as such.

Table A4: Areas for which none of the three indicators is positive (19 out of 41 on the September 13th 2023).

Province	WWTP	High	Fast	Incr.	Norm. viral load (%) ¹	Mean viral load ²	Norm evol. (%/week) ³	Incr. days ⁴	Date Max cc ⁵
Oost-Vlaanderen	Aalst	0	0	0	1.48	0.04	0	0	08/03/2023
Antwerpen	Aartselaar	0	0	0	3.8	0.02	0	0	08/03/2023
Antwerpen	Antwerpen-Noord	0	0	0	30.97	0.17	0	0	08/03/2023
Brabant Wallon	Basse Wavre (Dyle)	0	0	0	40.11	0.26	-39.1	0	15/02/2023
Vlaams-Brabant	Beersel	0	0	0	6.33	0.43	-45.5	0	9 th wave
West-Vlaanderen	Brugge	0	0	0	5.61	0.26	-48.8	0	04/01/2023
Brussels	Brussels-South	0	0	0	6.41	0.45	-29.4	0	03/10/2022

Province	WWTP	High	Fast	Incr.	Norm. viral load (%) ¹	Mean viral load ²	Norm. evol. (%/week) ³	Incr. days ⁴	Date Max cc ⁵
Antwerpen	Deurne	0	0	0	34.17	0.19	0	0	08/03/2023
Hainaut	Froyennes	0	0	0	24	0.27	-33.8	0	06/09/2023
Vlaams-Brabant	Grimbergen	0	0	0	0.51	0.07	0	0	9 th wave
West-Vlaanderen	Harelbeke	0	0	0	0.18	0.01	0	0	9 th wave
Limburg	Hasselt	0	0	0	0	0	0	0	08/03/2023
Hainaut	Marchienne-au-Pont	0	0	0	26.11	0.57	-39.6	0	9 th wave
Antwerpen	Mechelen-Noord	0	0	0	2.77	0.05	0	0	08/03/2023
Hainaut	Montignies-sur-Sambre	0	0	0	1.65	0.02	0	0	22/03/2023
West-Vlaanderen	Oostende	0	0	0	1.46	0.14	0	0	9 th wave
Hainaut	Roselies	0	0	0	47.85	0.48	-29.8	0	06/09/2023
Oost-Vlaanderen	Sint-Niklaas	0	0	0	18.79	0.48	-41.4	0	9 th wave
Antwerpen	Turnhout	0	0	0	3.31	0.08	0	0	01/03/2023

¹ : the viral load normalized with the maximum viral load measured in the corresponding catchment area during the ninth wave (i.e. from 21th of November 2022 till the first of January 2023).

² : the viral load computed on the mean of the replicates of the three targeted gene fragments as explained in section "3.2 Wastewater results". The mean viral load is expressed in 10¹² copies/day/100k inhabitants.

³ : the slope (%/week) of the past 7 days moving average of the viral load if the corresponding concentration is above the estimated limit of quantification.

⁴ : the cumulative number of days of increase of the past 14 days moving average of the viral load.

⁵ : date at which the measured viral load was the highest since the beginning of the ninth wave. If the date was between the 21th of November 2022 and the first of January 2023, the date is considered to be during the ninth wave and mentioned as such.

Table A5: Areas for which the data are missing on September 13th 2023.

Province	WWTP
Liège	Amay
Brussels	Brussels-North
Oost-Vlaanderen	Destelbergen
Limburg	Houthalen-Centrum
Liège	Liege (Grosses Battes)
Liège	Liege Oupeye
Liège	Liege Sclessin